

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
22 May 2003 (22.05.2003)

PCT

(10) International Publication Number  
**WO 03/041501 A1**

(51) International Patent Classification<sup>7</sup>: **A01N 43/40**,  
47/12

SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US,  
UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number: **PCT/EP02/13490**

(22) International Filing Date:

13 November 2002 (13.11.2002)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

01/14692

14 November 2001 (14.11.2001) **FR**

(71) Applicant (for all designated States except US): **BAYER CROPSCIENCE S.A.** [FR/FR]; 55, avenue René Cassin, F-69009 Lyon (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **WEGMANN, Thomas** [DE/FR]; 21, route de Saint-Romain, F-69450 Saint-Cyr-Au-Mont-d'Or (FR). **MERCER, Richard** [GB/FR]; 14, rue du Domaine, F-69130 Ecully (FR).

(74) Agent: **BAYER CROPSCIENCE S.A.**; Patents & Licensing Department, 14-20, rue Pierre Baizet, B.P. 9163, F-69263 Lyon (FR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW. ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only

**Published:**

- with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **FUNGICIDAL COMPOSITION BASED ON A PYRIDYLMETHYLBENZAMIDE DERIVATIVE AND A VALINAMIDE DERIVATIVE**

(57) Abstract: Fungicidal compositions comprising: a) at least one pyridylmethylbenzamide derivative of formula (I), in which the various radicals are as defined in the description; and b) at least one compound (II) of the valinamide type, preferably iprovalicarb or N<sup>1</sup>-[(R)-1-(6-fluoro-2-benzothiazolyl)ethyl]-N<sup>2</sup>-isopropoxycarbonyl-L-valinamide; and their agriculturally acceptable isomers and addition salts with an acid. Method for the curative or preventive control of phytopathogenic fungi of crops using an effective and non-phytotoxic quantity of one of these fungicidal compositions.

WO 03/041501 A1

**FUNGICIDAL COMPOSITION BASED ON A**  
**PYRIDYLMETHYLBENZAMIDE DERIVATIVE AND A**  
**VALINAMIDE DERIVATE**

5

The present invention relates to combinations of fungicidal compounds intended in particular for protecting crops against fungal diseases, and the corresponding methods of protection by application of the said combinations.

10 More precisely, the subject of the present invention is novel fungicidal compositions based on pyridylmethylbenzamide derivatives and derivatives of the valinamide type and in particular iprovalicarb having the chemical name 1-methylethyl[(1S)-2-methyl-1-[[[(1-(4-methylphenyl)ethyl]amino]carbonyl]carbamate or  $N^1$ -[(R)-1-(6-fluoro-2-benzothiazolyl)ethyl]- $N^2$ -isopropoxycarbonyl-L-valinamide.

15

As regards fungicidal activity, in particular for the protection of crops, one of the problems at the heart of the research studies carried out in this technical field is the improvement of performances, in particular in terms of fungicidal activity and in particular in terms of maintaining this fungicidal activity over time.

20

Naturally, the fungicidal compounds useful for the protection of plants against fungi must be endowed with an ecotoxicity which is reduced to the minimum. As far as possible, they should not be dangerous or toxic to the operator during use.

Furthermore, it is advantageous for fungicidal compounds to have a broad activity spectrum.

25

The economic factor should of course not be overlooked in the search for novel fungicidal compounds.

Without being limiting, attention is paid more particularly in the context of the invention to protection against infestation, by fungi, of grapevine, cereals, fruits, vegetables, lucerne, soyabean, market garden crops, turf, wood and horticultural plants, among others.

30

The composition according to the invention include one or more compounds of the pyridylmethylbenzamide type with fungicidal action as described in European patent application EP-A-1 056 723. These compounds make it possible to prevent the growth and development of phytopathogenic fungi (for example those which are active in the treatment of downy mildew, for example of grapevine, such as *Plasmopara viticola*).

35

This patent application EP-A-1 056 723 makes a general allusion to the possibility of combining these fungicidal active agents with products known to also develop a fungicidal activity, without citing by name examples of co-active ingredients, or even families of co-active ingredients, which are capable of being combined with pyridylmethylbenzamides.

One of the objectives of the present invention is to solve the problems set out above.

One of the objectives of the present invention is to provide novel fungicidal products which can be used, in particular by the farmer, for controlling the fungi infesting crops and in particular for controlling the major fungal diseases of grapevine such as downy mildew.

Another objective of the invention is to provide a novel fungicidal composition based on pyridylmethylbenzamide derivatives which is a lot more active against fungi which are harmful to plants, including grapevine, and which is in particular active over longer periods than the antifungal agents known up until now.

Two similar objectives of the present invention relate to potato crops.

Another objective of the invention is to provide a novel fungicidal agent which is completely high-performing in particular as regards its efficacy against fungi and the perennality of this efficacy so as to be able to reduce the doses of chemical products spread in the environment for combating fungal attacks of crops in particular vine downy mildew.

Another objective of the invention is to provide a novel fungicidal composition which is more active and active for longer, and which therefore has a lower dose, but which is also less toxic.

Another objective of the invention is to provide a novel broad-spectrum fungicidal composition which is perennially effective and which offers the farmer a large number of products so that the latter finds among them the product best suited to his particular use.

Another objective of the invention is to provide a novel fungicidal composition satisfying the specifications aimed at in the above objectives and which is also of a lower cost price, which is easy and which is not dangerous to handle.

Another objective of the invention is to provide a novel fungicidal composition as defined in the above objectives and which is useful in the preventive and curative treatment of fungal diseases, for example, of grapevine, cereals, Solanaceae, fruit, vegetables, lucerne, soyabean, market garden crops including potatoes, turf, wood or horticultural or ornamental plants.

Another objective of the invention is to provide a preventive and/or curative treatment of plants and in particular of crops, using a fungicidal composition or a fungicidal combination combining the products of the composition as defined in the above objectives, it being necessary for such a treatment to have a high and perennial efficacy against a very wide variety of fungi, while minimizing the doses, the toxicity and the cost.


Another objective of the present invention is to provide a kit for controlling, by way of curing and/or preventing and/or eradicating, the phytopathogenic fungi of plants and in particular of crops, such as grapevine, which meets the specifications set out in the objectives above.

An additional objective of the present invention is to allow improvement in the yield of the crops which is significant from an agronomic point of view.

All these objectives, among others, were achieved by the inventors who have had the merit of finding a fungicidal combination between pyridylmethylbenzamide derivatives and compounds of the valinamide type, preferably fungicides. Such a combination surprisingly and unexpectedly exhibiting a very high and perennial antifungal efficacy against a broad spectrum of fungi and in particular against those responsible for diseases such as vine downy mildew.

The present invention, which completely or partially satisfies the abovementioned objectives, therefore relates firstly to fungicidal compositions comprising:

a) at least one pyridylmethylbenzamide derivative of formula (I):



in which

- R¹ is chosen from a hydrogen atom, an optionally substituted alkyl radical and an optionally substituted acyl radical;

- R² is chosen from a hydrogen atom and an optionally substituted alkyl radical;

- R³ and R⁴, which are identical or different are independently chosen from a halogen atom, the hydroxyl radical, the cyano radical, the nitro radical, the

radical -SF<sub>5</sub>, the trialkylsilyl radical, an optionally substituted amino radical, an acyl radical, and a group E, OE or SE, in which E is chosen from an alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl and heterocyclyl radical, it being possible for each of them to be optionally substituted;

5                   • c represents 0, 1, 2, 3 or 4;

                  • q represents 0, 1, 2, 3 or 4;

                  and their agriculturally acceptable possible optical and/or geometric isomers, tautomers and addition salts with an acid or a base;

                  and

10               b) at least one compound (II) of the valinamide type, and their agriculturally acceptable isomers and addition salts with an acid.

In the definitions of the compounds of formula (I) set out above, the various radicals and chemical terms used have, unless otherwise stated, the following meanings:

15

                  • "alkyl or alkyl-" denotes a linear or branched saturated hydrocarbon radical containing from 1 to 6 carbon atoms;

                  • "alkenyl" denotes a linear or branched hydrocarbon radical containing from 2 to 6 carbon atoms and an unsaturation in the form of double bond;

20

                  • "alkynyl" denotes a linear or branched hydrocarbon radical containing from 2 to 6 carbon atoms and an unsaturation in the form of a triple bond;

                  • "alkoxy" denotes an alkyloxy radical;

                  • "acyl" denotes the formyl radical or an alkoxycarbonyl radical;

25

                  • "cycloalkyl" denotes a saturated cyclic hydrocarbon radical containing from 3 to 8 carbon atoms;

                  • "aryl" denotes a phenyl or naphthyl radical;

30

                  • "heterocyclyl" denotes an unsaturated or a completely or partially saturated cyclic radical containing from 3 to 8 atoms, chosen from carbon, nitrogen, sulphur and oxygen, for example, and without limitation, pyridyl, pyridinyl, quinolyl, furyl, thienyl, pyrrolyl, oxazolinyl;

                  • the term "optionally substituted " means that the radicals thus termed may be substituted with one or more radicals chosen from chlorine, bromine, fluorine, iodine, alkyl, alkoxy, hydroxyl, nitro, amino; cyano and acyl.

35

The compounds of formula (I) are, for example, described in patent application EP-A-1 056 723 and, among these, the following will be preferred:

compounds possessing one of the following characteristics:

•  $R^1$  and  $R^2$ , which are identical or different, are independently chosen from a hydrogen atom and an optionally substituted alkyl radical;

5      •  $R^3$  and  $R^4$ , which are identical or different, are independently chosen from a halogen atom, the hydroxyl radical, the nitro radical, an optionally substituted amino radical, an acyl radical, and a group E, OE or SE, in which E is chosen from an alkyl, cycloalkyl, phenyl and heterocyclyl radical, it being possible for each of them to be optionally substituted;

10      • c represents 0, 1, 2 or 3;

    • q represents 0, 1, 2 or 3;

and their agriculturally acceptable possible optical and/or geometric isomers, tautomers and addition salts with an acid or a base.

Among the compounds of formula (I), the following will also be preferred:

15      compounds possessing at least one of the following characteristics:

    •  $R^1$  and  $R^2$ , which are identical or different, are independently chosen from a hydrogen atom and a methyl or ethyl radical;

20      •  $R^3$  and  $R^4$ , which are identical or different, are independently chosen from a halogen atom, the nitro radical, an optionally substituted amino radical and an alkyl, cycloalkyl, phenyl or heterocyclyl radical, it being possible for each of them to be optionally substituted;

    • c represents 1 or 2;

    • q represents 1 or 2;

25      and their agriculturally acceptable possible optical and/or geometric isomers, tautomers and addition salts with an acid or a base.

More particularly, the compounds of formula (I) are:

products possessing the following characteristics:

    •  $R^1$  and  $R^2$  each represent a hydrogen atom;

30      •  $R^3$  and  $R^4$ , which are identical or different, are independently chosen from a halogen atom, the nitro radical, an alkyl radical and the trifluoromethyl radical;

    • c and q represent, independently of each other, 2;

and their agriculturally acceptable possible tautomers and addition salts with an acid or a base.

35      By way of examples, the following compounds of formula (I) are most particularly preferred in the context of the present invention:

- compound (Ia) which is 2,6-dichloro-N-{{3-chloro-5-(trifluoromethyl)-2-pyridinyl}methyl}benzamide;

- compound (Ib) which is N-{{3-chloro-5-(trifluoromethyl)-2-pyridinyl)methyl}}-2-fluoro-6-nitrobenzamide;

5                    • compound (Ic) which is N-{{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl}-2-methyl-6-nitrobenzamide,

and their agriculturally acceptable possible tautomers and addition salts with an acid or a base.

10           The compounds (II) are preferably of the valinamide type and are known for their fungicidal activity, and their agriculturally acceptable isomers and addition salts with an acid.

Still more preferably, the antifungal compound of the valinamide type (II) is chosen from the group of compounds comprising: iprovalicarb having the chemical name 1-methylethyl[(1S)-2-methyl-1-[[[(1-(4-methylphenyl)ethyl]amino)carbonyl]carbamate or  $N^1$ -(R)-1-(6-fluoro-2-benzothiazolyl)ethyl]- $N^2$ -isopropoxycarbonyl-L-valinamide.

Advantageously, the compositions according to the present invention comprise:  
20 compound (Ia) and/or compound (Ib) and/or compound (Ic) combined with  
iprovalicarb or with  $N^1$ -[(R)-1-(6-fluoro-2-benzothiazolyl)ethyl]- $N^2$ -isopropoxy-  
carbonyl-L-valinamide.

The preferred compositions of the present invention comprise compound (Ia) with iprovalicarb or with  $N^1$ -[(R)-1-(6-fluoro-2-benzothiazolyl)ethyl]- $N^2$ -isopropoxycarbonyl-L-valinamide, and the possible tautomers and addition salts with an acid or a base of compound (Ia) and of iprovalicarb or of  $N^1$ -[(R)-1-(6-fluoro-2-benzothiazolyl)ethyl]- $N^2$ -isopropoxycarbonyl-L-valinamide, as long as these equivalents are agriculturally acceptable.

The fungicidal combination of compounds (I) with compounds (II) according to the invention makes it possible to significantly improve the persistence of antifungal activity in the context of the curative and/or preventive treatment of major diseases of crops, including in particular those caused by fungi of the family of Peronosporaceae, in particular *Plasmopara viticola* (vine downy mildew), *Plasmopara halstedei* (sunflower mildew), *Pseudoperonospora* *sp* (in particular cucurbit mildew (*Pseudoperonospora cubensis*) and downy mildew of hops (*Pseudoperonospora humuli*)), *Bremia lactucae* (mildew of lettuce), *Peronospora tabacinae* (downy mildew of tobacco), *Peronospora*

*destructor* (downy mildew of onion), *Peronospora parasitica* (downy mildew of cabbage), *Peronospora farinosa* (downy mildew of chicory and downy mildew of beetroot).

5 In particular, this combination has eradicant properties which are superior to those of the products alone.

From the point of view of weight, it should be specified that in accordance with the invention, the compound (I)/compound (II) ratio ranges most often from 1/100 to 100/1, preferably from 1/10 to 10/1, more preferably from 1/5 to 5/1.

10 The compound (I)/compound (II) ratio is defined as being the ratio by weight of these 2 compounds. The same applies to any ratio of 2 chemical compounds, which is subsequently measured in the present text, since a definition different from this ratio is not expressly given.

15 According to another aspect of the present invention, in the compositions according to the invention, the compound (I)/compound (II) ratio may be advantageously chosen so as to produce a synergistic effect. The term synergistic effect is understood to mean in particular that defined by Colby in an article entitled "Calculation of the synergistic and antagonistic responses of herbicide combinations" Weeds, (1967), 15, pages 20-22.

20 The latter article mentions the formula:

$$E = x + y - \frac{x * y}{100}$$

25 in which E represents the expected percentage of inhibition of the disease for the combination of the two fungicides at defined doses (for example equal to x and y respectively), x is the percentage of inhibition observed for the disease by the compound (I) at a defined dose (equal to x), y is the percentage of inhibition observed for the disease by the compound (II) at a defined dose (equal to y). When the percentage of inhibition observed for the combination is greater than E, there is a synergistic effect.

30 The term "synergistic effect" also means the effect defined by application of the Tammes method, "Isoboles, a graphic representation of synergism in pesticides", Netherlands Journal of Plant Pathology, 70(1964), pages 73-80.

35 The compound (I)/compound (II) ratio ranges indicated above do not in any way limit the scope of the invention, but are, rather, mentioned as a guide, a person skilled in the art being entirely capable of carrying out additional tests to find other values of the ratio of doses of these two compounds, in particular for which a synergistic effect is observed.

Advantageously, the compound (I)/compound (II) ratio is chosen so as to produce a synergistic effect.

Usually, the compositions according to the invention comprise between 0.00001 and 100%, preferably between 0.001 and 80%, of active compounds, whether these compounds are combined, or whether they are in the form of two active ingredients used separately.

It is clearly understood that the said fungicidal compositions may contain a single compound (I) or more than such a compound and/or a single compound (II) or more than such as compound, and one or more other fungicidal, herbicidal, insecticidal and/or plant growth regulating compounds, according to the use for which they are intended.

Thus, the fungicidal compositions according to the present invention may also comprise, for example, one or more other fungicidal active ingredients chosen from acibenzolar-S-methyl, azoxystrobin, benalaxyl, benomyl, blasticidin-S, bromuconazole, captafol, captan, carbendazim, carboxin, carpropamid, chlorothalonil, the fungicidal compositions based on copper and copper derivatives such as copper hydroxide and copper oxychloride, cyazofamid, cyflufenamid, cymoxanil, cyproconazole, cyprodinyl, dichloran, diclocymet, diethofencarb, difenoconazole, diflumetorim, dimethomorph, dimoxystrobin, diniconazole, discostrobin, dodemorph, dodine, edifenphos, epoxyconazole, ethaboxam, ethirimol, fenarimol, fenbuconazole, fenhexamid, fenoxanil, fenpiclonil, fenpropidin, fenpropimorph, ferimzone, fluazinam, fludioxonil, flumetover, fluquinconazole, flusilazole, flusulfamide, flutolanil, flutriafol, folpel, furalaxyl, furametpyr, guazatine, hexaconazole, hymexazol, imazalil, iprobenphos, iprodione, isoprothiolane, kasugamycin, kresoxim-methyl, mefenoxam, mepanipyrim, metalaxyl and its entantiomeric forms such as metalaxyl-M, metconazole, metiram-zinc, metominostrobin, metrafenone, nicobifen, oxadixyl, oxpoconazole, pefurazoate, penconazole, pencycuron, phosphorous acid and its derivatives such as fosetyl-Al, phthalide, picoxystrobin, probenazole, prochloraz, procymidone, propamocarb, propiconazole, pyraclostrobin, pyrimethanil, pyroquilon, quinoxifen, silthiofam, simeconazole, spiroxamine, sup-[190, tebuconazole, tetraconazole, thiabendazole, thifluzamide, thiophanate, for example thiophanate-methyl, thiram, tiadinil, triadimefon, triadimenol, tricyclazole, tridemorph, trifloxystrobin, triticonazole, vinclozolin and zoxamide.

In addition to these additional active agents, the fungicidal compositions according to the invention may also contain any other excipient and/or auxiliary agent

useful in plant protection formulations such as, for example, an agriculturally suitable inert carrier and optionally an agriculturally suitable surfactant.

As regards the presentations of the compositions according to the invention, it should be indicated that they are appropriate for a large number of formulations. Thus, it is possible to use these compositions as aerosol dispenser; bait (ready-to-use); concentrate for preparation of baits; stock bait; suspension of capsules; cold fogging concentrate; dustable powder; emulsifiable concentrate; aqueous/aqueous type emulsion; oil/inverse type emulsion; encapsulated granule; fine granule; suspension concentrate for seed treatment; compressed gas; gas generating product; grain bait; granular bait; granule; hot fogging concentrate; macrogranule; microgranule; oil-dispersible powder, oil miscible suspension concentrate; oil-miscible liquid; paste; plant rodlet; plate bait; powder for dry seed treatment; scrap bait; seeds coated with a pesticide; smoke candle; smoke cartridge; smoke generator; smoke pellet; smoke rodlet; smoke tablet; smoke tin; soluble concentrate; soluble powder; solution for seed treatment; suspension concentrate (= flowable concentrate); tracking powder; ultra low volume liquid; ultra low volume suspension; vapour releasing product; water-dispersible granules or tablets; water dispersible powder for slurry treatment; water-soluble granules or tablets; water-soluble powder for seed treatment; wettable powder.

These compositions cover not only the compositions which are ready to be applied to the crop to be treated by means of a suitable device, such as a spraying device, but also the commercial concentrated compositions which have to be diluted before application to the crop.

The compositions described below are used in general for application to growing plants, or to sites where crops are grown, or for the coating or film-coating of seeds.

The compositions according to the invention are, appropriately, applied to the vegetation and in particular to the leaves infested or capable of being infested with the phytopathogenic fungi. Another method of applying the compounds or compositions according to the invention is to add a formulation containing the active ingredients to the irrigation water. This irrigation may be an irrigation using sprinklers.

For their use in practice, the compositions according to the invention can be used alone and can also advantageously be used in formulations containing one or the other of the active ingredients or alternatively both of them together, in combination or association with one or more other compatible components which are, for example, solid or liquid fillers or diluents, adjuvants, surfactants or equivalents, which are suitable for the desired use and which are acceptable for uses in agriculture. The formulations can be of any type known in the sector which are suitable for application

onto all types of plantations or crops. These formulations, which can be prepared in any manner known in this sector, also form part of the invention.

The formulations can also contain ingredients of other types, such as protective colloids, adhesives, thickeners, thixotropic agents, penetrating agents, oils for spraying, stabilizers, preserving agents (in particular mouldproofing agents), sequestering agents or the like, as well as other known active ingredients which have pesticidal properties (in particular fungicidal, insecticidal, acaricidal or nematocidal properties) or which have properties of regulating plant growth. More generally, the compounds used in the invention can be combined with any solid or liquid additives corresponding to the usual formulation techniques.

In the present account, the term "filler" means an organic or inorganic, natural or synthetic component with which the active components are combined to facilitate its application, for example, onto the plants, the seeds or the soil. This filler is consequently generally inert and it must be acceptable (for example acceptable for agronomic uses, in particular for treating plants).

The filler can be solid, for example clays, natural or synthetic silicates, silica, resins, waxes, solid fertilizers (for example ammonium salts), natural soil minerals, such as kaolins, clays, talc, lime, quartz, attapulgite, montmorillonite, bentonite or diatomaceous earths, or synthetic minerals, such as silica, alumina or silicates, in particular aluminium or magnesium silicates. The solid fillers which are suitable for granules are as follows: natural, crushed or broken rocks, such as calcite, marble, pumice, sepiolite and dolomite; synthetic granules of inorganic or organic flours; granules of organic material such as sawdust, coconut shell, corn ear or envelope, or tobacco stem; kieselguhr, tricalcium phosphate, powdered cork or adsorbent carbon black; water-soluble polymers, resins, waxes; or solid fertilizers. Such compositions can, if so desired, contain one or more compatible agents such as wetting agents, dispersing agents, emulsifiers or colourings which, when they are solid, can also act as diluents.

The fillers can also be liquid, for example: water, alcohols, in particular butanol or glycol, as well as ethers or esters thereof, in particular methyl glycol acetate; ketones, in particular acetone, cyclohexanone, methyl ethyl ketone, methyl isobutyl ketone or isophorone; petroleum fractions such as paraffinic or aromatic hydrocarbons, in particular xylenes or alkylnaphthalenes; mineral or plant oils; aliphatic chlorohydrocarbons, in particular trichloroethane or methylene chloride; aromatic chlorohydrocarbons, in particular chlorobenzenes; water-soluble or highly polar solvents such as dimethylformamide, dimethyl sulfoxide, N,N-dimethyl-acetamide or N-

methylpyrrolidone; N-octylpyrrolidone, liquefied gases; or the like, whether they are taken separately or as a mixture.

5 The surfactant can be an emulsifier, a dispersing agent or a wetting agent, of ionic or nonionic type or a mixture of these surfactants. Among those surfactants there are used, for example, polyacrylic acid salts, lignosulphonic acid salts, phenolsulphonic or naphthalenesulphonic acid salts, polycondensates of ethylene oxide with fatty  
10 alcohols or fatty acids or fatty esters or fatty amines, substituted phenols (in particular alkylphenols or arylphenols), ester-salts of sulphosuccinic acid, taurine derivatives (in particular alkyl taurates), phosphoric esters of alcohols or of polycondensates of ethylene oxide with phenols, fatty acid esters with polyols, or sulphate, sulphonate or phosphate functional derivatives of the compounds described above. The presence of at least one surfactant is generally essential when the active ingredients and/or the inert  
15 filler are insoluble or only sparingly soluble in water and when the filler for the said composition to be applied is water.

20 The formulations according to the invention can also contain other additives such as adhesives or colourings. Adhesives such as carboxymethylcellulose, or natural or synthetic polymers in the form of powders, granules or matrices, such as gum arabic, latex, polyvinylpyrrolidone, polyvinyl alcohol or polyvinyl acetate, natural phospholipids, such as cephalins or lecithins, or synthetic phospholipids can be used in the formulations. It is possible to use colourings such as inorganic pigments, such as, for example: iron oxides, titanium oxides, Prussian blue; organic colouringstuffs, such as those of the alizarin, azo or metal phthalocyanin type; or of trace elements such as iron, manganese, boron, copper, cobalt, molybdenum or zinc salts.

25 The formulations containing the compositions of the invention, which are used to control the phytopathogenic fungi of crops, can also contain stabilizers, other fungicidal agents, insecticides, acaricides, nematocides, anti-helminths or anti-coccidoses, bactericides, attractant or repellent agents, deodorizers, flavourings or colourings.

30 These can be chosen for the purpose of improving the strength, the persistence, the safety, and the spectrum of action on the phytopathogenic fungi of crops or to make the composition capable of accomplishing other useful functions for the areas treated.

As regards the preparation of compounds (I), reference may be made to European patent application EP-A-1 056 723.

35 As regards the production of compounds (II), reference may be made to the book "The Electronic Pesticide Manual – Version 1.0" – British Crop Protection Council – Ed Clive Tomlin or to European patent application No. 775 696.

According to another of these objects, the invention relates to a method for controlling, by way of curing, preventing or eradicating, the phytopathogenic fungi of crops, characterized in that an effective (agronomically effective) and nonphytotoxic quantity of a fungicidal composition as defined above is applied to the soil where plants grow or are capable of growing, to the leaves and/or the fruits of plants or to the seeds of plants.

In this method, a composition is used which is prepared beforehand by mixing the 2 active compounds (I) and (II).

According to a variant of such a method of controlling, by way of curing, preventing or eradicating, the phytopathogenic fungi of crops:

- a combination of at least one compound (I) and of at least one compound (II) as defined above is used;

- the compounds (I) and (II) are applied simultaneously, separately or sequentially to the soil where plants grow or are capable of growing, to the leaves and/or the fruits of plants or to the seeds of plants, an effective (agronomically effective) and nonphytotoxic quantity.

This variant corresponds to a fresh preparation of the fungicidal composition.

It is also possible to apply simultaneously, successively or separately so as to have the conjugated (I)/(II) effect, of a composition each containing one of the two active ingredients (I) or (II).

Preferably, the fungicidal compositions according to the invention usually contain from 0.5 to 95% of the combination of compound (I) and compound (II). This may be the concentrated composition, that is to say the commercial product combining compound (I) and compound (II). This may also be the dilute composition ready to be applied to the crops to be treated. In the latter case, the dilution with water may be carried out either using a commercial concentrated composition containing compound (I) and compound (II) (this mixture is called ready mix), or using the tank mix of two commercial concentrated compositions each containing compound (I) and compound (II).

The treatment of crops against phytopathogenic diseases, using the fungicidal composition according to the invention, is carried out, for example, by application or by administration, with an effective and nonphytotoxic quantity of the abovementioned fungicidal composition or combination, to the aerial parts of the crops or to the soil where they grow, the said crops being those which are infested or which are capable of being infested by a phytopathogenic disease such as downy mildew, oidium, brown rust or Septoria disease. The expression treatment of the crop is also understood to mean the

treatment of the reproductive products of the crop, such as the seeds or the tubers for example.

Under specific conditions, for example according to the nature of the phytopathogenic fungus to be treated, a lower dose may offer adequate protection. Conversely, certain climatic conditions, resistance or other factors may require higher doses of active ingredient.

The effective working doses of the combinations used in the invention can vary within wide proportions, in particular depending on the nature of the phytopathogenic fungi to be eliminated or the degree of infestation, for example, of the plants with these fungi.

The optimum dose usually depends on several factors, for example on the type of phytopathogenic fungus to be treated, on the type or level of development of the infested plant, on the density of vegetation, or alternatively on the method of application. More preferably, an effective dose of active ingredients (I) and (II) is between about 0.1 g/ha and about 1 000 g/ha.

Without it being limiting, the crop treated with the fungicidal composition or combination according to the invention is, for example, a cereal, but this could be grapevine, vegetables, fruits, lucerne, soyabean, market garden crops, turf, wood or horticultural plants.

The phytopathogenic fungi of crops which may be controlled by this method are selected from the group comprising:

- the group of oomycetes:

- of the family of Peronosporaceae, in particular *Plasmopara viticola* (vine downy mildew), *Plasmopara halstedei* (sunflower mildew), *Pseudoperonospora sp* (in particular cucurbit mildew (*Pseudoperonospora cubensis*) and downy mildew of hops (*Pseudoperonospora humuli*)), *Bremia lactucae* (mildew of lettuce), *Peronospora tabacinae* (downy mildew of tobacco), *Peronospora destructor* (downy mildew of onion), *Peronospora parasitica* (downy mildew of cabbage), *Peronospora farinosa* (downy mildew of chicory and downy mildew of beetroot);

- of the genus *Phytophthora* such as *Phytophthora phaseoli*, *Phytophthora citrophthora*, *Phytophthora capsici*, *Phytophthora cactorum*, *Phytophthora palmivora*, *Phytophthora cinnamoni*, *Phytophthora megasperma*, *Phytophthora parasitica*, *Phytophthora fragariae*, *Phytophthora cryptogea*, *Phytophthora porri*, *Phytophthora nicotianae*, *Phytophthora infestans* (mildew of Solanaceae, in particular late blight of potato or tomato);

- the group of adelomycetes (ascomycetes):

- of the genus *Alternaria*, for example *Alternaria solani* (early blight of Solanaceae and in particular of tomato and potato),
- of the genus *Guignardia*, in particular *Guignardia bidwelli* (black rot of grapevine),
- 5 - of the genus *Venturia*, for example *Venturia inaequalis*, *Venturia pirina* (apple or pear scabs),
- of the genus *Oidium*, for example powdery mildew of grapevine (*Uncinula necator*); oidium of leguminous crops, for example *Erysiphe polygoni* (powdery mildew of Cruciferae); *Leveillula taurica*, *Erysiphe cichoracearum*, *Sphaerotheca fuliginea*
- 10 (powdery mildew of cucurbits, of composites and of tomato); *Erysiphe communis* (powdery mildew of beetroot and cabbage); *Erysiphe pisi* (powdery mildew of pea and lucerne); *Erysiphe polyphaga* (powdery mildew of haricot bean and cucumber); *Erysiphe umbelliferarum* (powdery mildew of umbellifera, in particular of carrot); *Sphaerotheca humuli* (hop mildew); powdery mildew of wheat and barley (*Erysiphe graminis forma specie tritici* and *Erysiphe graminis forma specie hordei*),
- 15 - of the genus *Taphrina*, for example *Taphrina deformans* (peach leaf curl),
- of the genus *Septoria*, for example *Septoria nodorum* or *Septoria tritici* (*Septoria* disease of cereals),
- of the genus *Sclerotinia*, for example *Sclerotinia sclerotinium*,
- 20 - of the genus *Pseudocercospora*, for example *P. herpotrichoides* (eyespot of cereals),
- of the genus *Botrytis cinerea* (grapevine, vegetable and market garden crops, pea and the like),
- of the genus *Phomopsis viticola* (excoriosis of grapevine),
- 25 - of the genus *Pyrenospora*,
- of the genus *Helminthosporium*, for example *Helminthosporium tritici repentis* (yellow leaf spot of wheat) or *Helminthosporium teres* (yellow leaf spot of barley),
- of the genus *Drechslera* or *Pyrenophora*,
- of the group of basidiomycetes :
- 30 - of the genus *Puccinia*, for example *Puccinia recondita* or *striiformis* (wheat rust), *Puccinia triticina*, *Puccinia hordei*,
- of the family *Rhizoctonia* spp, for example *Rhizoctonia solani*.

35 In addition to their fungicidal activities at the heart of the invention, the compositions or combinations defined above may also have a biocide action against bacteria and viruses, such as for example:

- fire blight, *Erwinia amylovora*;
- bacterial streak of stone fruit trees, *Xanthomonas campestris*;
- pear blossom blight, *Pseudomonas syringae*;
- bacteriosis of rice and cereals;
- 5 - the viruses present on rice, vegetable and cereal crops.

The crops envisaged in the context of the present invention are preferably grapevine, cereal crops (wheat, barley, maize, rice) and vegetable crops (haricot bean, onion, cucurbitaceae, cabbage, potato, tomato, sweet pepper, spinach, pea, lettuce, 10 celery, chicory), fruit crops (strawberry plants, raspberry plants), tree crops (apple trees, pear trees, cherry trees, ginseng, lemon trees, coconut palms, pecan trees, cacao trees, walnut trees, rubber trees, olive trees, poplars, banana trees), sunflower, beetroot, tobacco, hop and ornamental crops.

A classification made, no longer based on the fungi or bacteria targeted, but on 15 the target crops may be illustrated as below:

- grapevine: downy mildew (*Plasmopara viticola*), powdery mildew (*Uncinula necator*), grey mould (*Botrytis cinerea*), excoriosis (*Phomopsis viticola*) and black rot (*Guignardia bidwelli*);
- Solanaceae: blight (*Phytophthora infestans*), alternara disease (*Alternaria solani*) and 20 grey mould (*Botrytis cinerea*),
- vegetable crops: downy mildew (*Peronospora* sp., *Bremia lactucae*, *Pseudoperonospora* sp), alternara (*Alternaria* sp.), sclerotinia disease (*Sclerotinia* sp.), grey mould (*Botrytis cinerea*), foot or root rot (*Rhizoctonia* spp.), powdery mildew (*Erysiphe* sp.; *Sphaerotheca fuliginea*),
- 25 -arboriculture: scab (*Venturia inaequalis*, *V. pirina*), bacterial diseases (*erwinia amylovora*, *xanthomonas campestris*, *pseudomonas syringae*), powdery mildew (*Podosphaera leucotricha*) and Monilia (*Monilia fructigena*),
- citrus: scab (*Elsinoe fawcetti*), melanose (*Phomopsis citri*) and *Phytophthora* sp. diseases,
- 30 - wheat, as regards controlling the following seed diseases: Fusarium diseases (*Microdochium nivale* and *Fusarium roseum*), smuts (*Tilletia caries*, *Tilletia controversa* or *Tilletia indica*), Septoria disease (*Septoria nodorum*),
- wheat, as regards controlling the following diseases of the aerial parts of the plant: eyespot (*Pseudocercospora herpotrichoïdes*), take-all (*Gaeumannomyces graminis*),
- 35 Fusarium disease of the foot (*F. culmorum*, *F. graminearum*), Rhizoctonia disease (*Rhizoctonia cerealis*), powdery mildew (*Erysiphe graminis forma specie tritici*), rusts

- (*Puccinia striiformis* and *Puccinia recondita*), Septoria diseases (*Septoria tritici* and *Septoria nodorum*) and yellow leaf spot of wheat (*Helminthosporium tritici-vulgaris*);
- wheat and barley, as regards controlling bacterial and viral diseases, for example barley yellow mosaic;
  - 5 - barley, as regards controlling the following seed diseases: yellow leaf spot (*Pyrenophora graminea*, *Bipolaris*, *Pyrenophora teres* and *Cochliobolus sativus*), loose smut (*Ustilago nuda*) and Fusarium diseases (*Microdochium nivale* and *Fusarium roseum*);
  - barley, as regards controlling the following diseases of the aerial parts of the plant:  
10 eyespot (*Pseudocercospora herpotrichoides*), yellow leaf spot (*Pyrenophora teres* and *Cochliobolus sativus*), powdery mildew (*Erysiphe graminis* forma specie hordei), dwarf leaf rust (*Puccinia hordei*) and leaf blotch (*Rhynchosporium secalis*);
  - potato, as regards controlling tuber diseases (in particular *Helminthosporium solani*, *Phoma tuberosa*, *Rhizoctonia solani*, *Fusarium solani*) and certain virus diseases  
15 (virus Y);
  - cotton, as regards controlling the following diseases of young plants obtained from seeds: damping-off diseases and collar rot (*Rhizoctonia solani*, *Fusarium oxysporum*), black root rot (*Thielaviopsis basicola*);
  - pea, as regards controlling the following seed diseases: anthracnose (*Ascochyta pisi*,  
20 *Mycosphaerella pinodes*), Fusarium disease (*Fusarium oxysporum*), grey mould (*Botrytis cinerea*), rust (*Uromyces pisi*);
  - rape plant, as regards controlling the following seed diseases: *Phoma lingam* and *Alternaria brassicae*, grey mould (*Botrytis cinerea*), and sclerotinia disease (*Sclerotinia sclerotinium*);
  - 25 - maize, as regards controlling seed diseases (*Rhizopus* sp., *Penicillium* sp., *Trichoderma* sp., *Aspergillus* sp. and *Gibberella fujikuroi*), yellow leaf spot (*Bipolaris*), Fusarium disease (*Fusarium oxysporum*);
  - rice: foot and root rot (*Rhizoctonia* spp.);
  - flax, as regards controlling seed disease (*Alternaria linicola*);
  - 30 - banana: *Cercospora* disease (*Mycosphaerella figiensis*);
  - turf: rust, powdery mildew, yellow leaf spot, terruric diseases (*Microdochium nivale*, *Pythium* sp., *Rhizoctonia solani*, *Sclerotinia homeocarpa*);
  - forest trees, as regards controlling damping-off (*Fusarium oxysporum*, *Rhizoctonia solani*).

Very advantageously, the method for controlling plant diseases according to the invention has shown excellent results against grapevine diseases: downy mildew (*Plasmopara viticola*), powdery mildew (*Uncinula necator*), grey mould (*Botrytis cinerea*), excoriosis (*Phomopsis viticola*) and black rot (*Guignardia bidwelli*) and against vegetable crop diseases.

The expression "are applied to the plants to be treated" is understood to mean, for the purposes of the present text, that the fungicidal compositions which are the subject of the invention may be applied by means of various methods of treatment such as:

- spraying onto the aerial parts of the said plants a liquid comprising one of the said compositions,
- dusting, the incorporation into the soil of granules or powders, spraying, around the said plants, and in the case of trees injection or daubing,
- coating or film-coating the seeds of the said plants with the aid of a plant-protection mixture comprising one of the said compositions.

Spraying a liquid onto the aerial parts of the crops to be treated is the preferred method of treatment.

The subject of the present invention is also a product comprising a compound (I) and a compound (II) as a combined preparation for simultaneous, separate or sequential use in controlling the phytopathogenic fungi of crops at a site.

Another object of the invention which is linked to the mode of preparing the composition according to the invention immediately before use consists of a kit for controlling, curatively or preventively, the phytopathogenic fungi of crops, characterized in that it comprises at least one compound (I) and at least one compound (II) as defined above, intended to be combined or used simultaneously, separately or sequentially in controlling the phytopathogenic fungi of crops at a site.

It is therefore a pack in which the user finds all the ingredients for preparing the fungicidal formulation which they wish to apply to the crops. These ingredients, which comprise in particular the active agents (I) and (II) and which are packaged separately, are provided, for example, in the form of a powder or in the form of a liquid which is concentrated to a greater or lesser degree. The user simply has to mix in the prescribed doses and to add the quantities of liquid, for example of water, necessary to obtain a formulation which is ready to use and which can be applied to the crops.

Most appropriate is a product for simultaneous, separate, alternate or sequential application of at least one fungicidal compound (I) and one fungicidal compound (II).

It goes without saying that the different variants which may be envisaged of the compositions and methods according to the invention form an integral part of the present invention; it being possible for the said different variants, moreover, to be combined or associated with each other without as a result departing either from the spirit or from the scope of the said invention.

In the same manner, the different aspects of the present invention which have just been described may be combined or associated with each other without as a result departing either from the spirit or from the scope of the said invention.

The following example is given purely by way of illustration of the invention and does not limit it in any manner.

**Example of treatment of vine cuttings and of potato tubers with a composition according to the invention**

This example demonstrates a synergy between a compound A of formula (I) and a compound B of formula (II) according to the present invention on vine downy mildew (*Plasmopara viticola*) and potato blight (*Phytophthora infestans*).

Starting with 2,6-dichloro-N-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl]benzamide (compound A) in the form of water-dispersible granules (WG) at 97.9 g/l and iprovalicarb (compound B) in the form of a suspension concentrate (SC) at 500 g/l, plant protection mixtures for application corresponding to a volume of 300 l of liquid per ha are prepared.

These two liquids are mixed so as to study a compound A/compound B ratio equal to 1/0, 1/5, 1/1, 5/1 and 0/1 for potato and 1/0, 1/5, 1/1 and 0/1 for grapevine.

For each ratio, the dose of compound A is fixed and consequently the dose range of compound B is varied.

Vine cuttings (*Vitis vinifera*), Cabernet sauvignon variety, and potato tubers (*Solanum tuberosum*), Bintje variety, are cultivated in pots.

For grapevine, when these plants are 2 months old (8- to 10-leaf stage, height 10 to 15 cm), they are treated by spraying a solution comprising compound A, or a solution comprising compound B, or alternatively a solution comprising compound A and compound B mixed in one of the ratios indicated above. The dose range corresponds to 100 mg/l, 50 mg/l, 20 mg/l, 10 mg/l and 1 mg/l of compound if formula (Ia).

When the potato plants reach the age of 5 weeks (7- to 8- leaf stage, height 10 to 40 cm), they undergo the same treatment with the following dose range: 100 mg/l, 50 mg/l, 10 mg/l, 5 mg/l and 1 mg/l of compound A.

Plants used as controls are treated with water containing no active ingredients.

5

After 48 h, the infection of the vine leaves and of the potato tubers is carried out by spraying an aqueous suspension adjusted to 100 000 sporocysts/ml of *Plasmopara viticola* onto the vine plants and an aqueous suspension adjusted to 40 000 sporocysts/ml of *Phytophthora infestans* onto the potato plants.

10

The infected plants are then incubated at 20°C and 90% relative humidity for 8 days for grapevine and at 16°C and 90% relative humidity for 6 days for potato.

15

The results are then observed, in comparison with the untreated but also infected control plants. Further, the infected surface area is estimated on all the leaves of a plant, compared with the controls. The level of infection is thus obtained and the efficacy of the treatment, expressed as a percentage (%) is calculated according to the Abbott formula.

20

The results obtained are presented in the form of points, corresponding to 50, 70 or 90% destruction of the parasite and placed in a TAMMES isobole diagram which comprises on the x-axis the doses of compound A expressed in mg/l and on the y-axis the doses of compound B also in mg/l.

### **Results :**

#### **1) 48 h preventive application on vine downy mildew**

25

The calculated results corresponding to the ED50 and 70 values (effective doses providing 50 to 70% control of the disease respectively) for the different ratios of the mixtures studied, calculated on the basis of 3 repeats per factor, are presented in Table 1. Also presented in this table are the limits of the confidence interval in which these ED values exist for a 95% risk.

	ED50 (confidence interval)	ED70 (confidence interval)
Compound A	6,6 ( 4 - 10,9 )	12,5 ( 8 - 19,6 )
Compound B	7,3 ( 4,9 - 11 )	17,5 ( 11,8 - 25,9 )
Compound A + Compound	2,3 ( 1,2 - 4,5 )	5,8 ( 3,4 - 10,1 )

<b>B (A/B ratio = 1)</b>		
<b>Compound A + Compound B (A/B ratio = 1/5)</b>	0,3 ( 0,1 - 1,1 )	1,3 ( 0,6 - 2,9 )

**Table 1: ED50 and ED70 following a preventive application of the mixture of compound A and of compound B onto vine downy mildew**

5           The results of this Table 1 are taken up in Figures 1 and 2 illustrating the study of synergy according to the TAMMES method for the ED50 and ED70 values, respectively.

10           The diagram of Figure 1 shows that the composition compound A + compound B for A/B ratios of 1:5 and 1:1 is in the zone of synergy according to the TAMMES method.

          Likewise, the diagram of Figure 2 shows that the composition compound A + compound B for A/B ratios of 1:5 and 1:1 is in the zone of synergy according to the TAMMES method.

15           **2) 48 h preventive application on potato blight**

          The calculated results corresponding to the ED50, 70 and 90 values (effective doses providing 50, 70 and 90% control of the disease respectively) for the different ratios of the mixtures studied, calculated on the basis of 3 repeats per factor, are presented in Table 2. Also presented in this table are the limits of the confidence interval in which these ED values exist for a 95% risk.

20

	ED50 (confidence interval)	ED70 (confidence interval)	ED90 (confidence interval)
<b>Compound A</b>	19,9 ( 12 - 33 )	43,5 ( 24,5 - 77,2 )	151,2 ( 61,5 - 371,8 )
<b>Compound B</b>	27,8 ( 18,3 - 42,2 )	69,3 ( 43,7 - 109,9 )	296,6 ( 136,2 - 646 )
<b>Compound A + Compound B (A/B ratio = 1)</b>	5,2 ( 3 - 9,2 )	14,9 ( 9 - 24,7 )	79,2 ( 34,8 - 180,5 )
<b>Compound A + Compound B (A/B ratio = 5/1)</b>	7,6 ( 4,3 - 13,5 )	19,2 ( 10,6 - 34,9 )	84,1 ( 33,4 - 211,9 )
<b>Compound A + Compound B (A/B ratio = 1/5)</b>	2,2 ( 1,2 - 4 )	5,4 ( 3,3 - 8,9 )	23,1 ( 12 - 44,5 )

**Table 2: ED50, ED70 and ED90 following a preventive application of the mixture of compound A and of compound B onto potato blight**

5

The results of this Table 2 are taken up in Figures 3 and 5 illustrating the study of synergy according to the TAMMES method for ED50, ED70 and ED90 values, respectively.

10


The diagram of Figure 3 shows that the composition compound A + compound B for A/B ratios of 1:5, 1:1 and 5:1 is in the zone of synergy according to the TAMMES method.

Likewise, the diagrams of Figures 4 and 5 show a synergy of the composition compound A + compound B for A/B ratios of 1:5, 1:1 and 5:1.

CLAIMS

1. Fungicidal compositions comprising:

5 a) at least one pyridylmethylbenzamide derivative of formula (I):



in which

10 • R¹ is chosen from a hydrogen atom, an optionally substituted alkyl radical and an optionally substituted acyl radical;

• R² is chosen from a hydrogen atom and an optionally substituted alkyl radical;

15 • R³ and R⁴, which are identical or different are independently chosen from a halogen atom, the hydroxyl radical, the cyano radical, the nitro radical, the radical -SF₅, the trialkylsilyl radical, an optionally substituted amino radical, an acyl radical, and a group E, OE or SE, in which E is chosen from an alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl and heterocyclyl radical, it being possible for each of them to be optionally substituted;

20 • c represents 0, 1, 2, 3 or 4;

• q represents 0, 1, 2, 3 or 4;

and their agriculturally acceptable possible optical and/or geometric isomers, tautomers and addition salts with an acid or a base;

and

25 b) at least one compound (II) of the valinamide type, and their agriculturally acceptable isomers and addition salts with an acid.

2. Fungicidal compositions according to Claim 1, characterized in that compound (I) possesses one of the following characteristics, taken in isolation or in combination:

30 • R¹ and R², which are identical or different, are independently chosen from a hydrogen atom and an optionally substituted alkyl radical;

•  $R^3$  and  $R^4$ , which are identical or different, are independently chosen from a halogen atom, the hydroxyl radical, the nitro radical, an optionally substituted amino radical, an acyl radical, and a group E, OE or SE, in which E is chosen from an alkyl, cycloalkyl, phenyl and heterocyclyl radical, it being possible for each of them to be optionally substituted;

• c represents 0, 1, 2 or 3;

• q represents 0, 1, 2 or 3;

and their agriculturally acceptable possible optical and/or geometric isomers, tautomers and addition salts with an acid or a base.

3. Fungicidal compositions according to Claim 1, characterized in that compound (I) possesses one of the following characteristics, taken in isolation or in combination:

•  $R^1$  and  $R^2$ , which are identical or different, are independently chosen from a hydrogen atom and a methyl or ethyl radical;

•  $R^3$  and  $R^4$ , which are identical or different, are independently chosen from a halogen atom, the nitro radical, an optionally substituted amino radical and an alkyl, cycloalkyl, phenyl or heterocyclyl radical, it being possible for each of them to be optionally substituted;

• c represents 1 or 2;

• q represents 1 or 2;

and their agriculturally acceptable possible optical and/or geometric isomers, tautomers and addition salts with an acid or a base.

4. Fungicidal compositions according to Claim 1, characterized in that compound (I) possesses one of the following characteristics:

•  $R^1$  and  $R^2$  each represent a hydrogen atom;

•  $R^3$  and  $R^4$ , which are identical or different, are independently chosen from a halogen atom (preferably Cl), the nitro radical, an alkyl radical and the trifluoromethyl radical;

• c and q represent, independently of each other, 2;

and their agriculturally acceptable possible optical and/or geometric isomers, tautomers and addition salts with an acid or a base.

5. Fungicidal compositions according to Claim 1, characterized in that the compound of formula (I) is chosen from:

• 2,6-dichloro-N-{{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl}-benzamide;

• N-{{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl}-2-fluoro-6-nitrobenzamide; and

5       • N-{{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl}-2-methyl-6-nitrobenzamide,

and their agriculturally acceptable possible tautomers and addition salts with an acid or a base.

10       6. Fungicidal compositions according to Claim 1, characterized in that compound (II) is of the valinamide type with fungicidal activity, and their agriculturally acceptable isomers and addition salts with an acid.

15       7. Fungicidal compositions according to Claim 6, characterized in that compound (II) is selected from the group comprising: iprovalicarb having the chemical name 1-methylethyl[(1S)-2-methyl-1-[[[1-(4-methylphenyl)ethyl]amino]carbonyl]carbamate or *N*<sup>1</sup>-(R)-1-(6-fluoro-2-benzothiazolyl)ethyl]-*N*<sup>2</sup>-isopropoxycarbonyl-L-valinamide.

20       8. Fungicidal compositions according to Claim 5 and Claim 7.

25       9. Fungicidal compositions according to Claim 8, characterized in that compound (I) is 2,6-dichloro-N-{{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl}benzamide and in that compound (II) is iprovalicarb or *N*<sup>1</sup>-(R)-1-(6-fluoro-2-benzothiazolyl)ethyl]-*N*<sup>2</sup>-isopropoxycarbonyl-L-valinamide.

30       10. Fungicidal compositions according to one of the preceding claims, characterized in that they comprise at least one compound of formula (I) and at least one compound (II), the compound (I)/compound (II) ratio ranging from 1/100 to 100/1, preferably from 1/10 to 10/1, still more preferably ranging from 1/5 to 5/1.

35       11. Fungicidal compositions according to Claim 10, characterized in that the compound (I)/compound (II) ratio is chosen so as to produce a synergistic effect.

12. Synergistic fungicidal compositions according to the preceding claim, characterized in that the compound (I)/compound (II) ratio is between 1/3 and 1/2.

13. Fungicidal compositions according to one of the preceding claims, characterized in that they comprise, in addition to compounds (I) and (II), an agriculturally suitable inert carrier and optionally an agriculturally suitable surfactant.
- 5 14. Fungicidal compositions according to one of the preceding claims, characterized in that they comprise between 0.00001 and 100%, preferably between 0.001 and 80%, of the combination of compound (I) and of compound (II).
- 10 15. Method for the curative or preventive control of phytopathogenic fungi of crops, characterized in that an effective (agronomically effective) and non-phytotoxic quantity of a fungicidal composition according to one of Claims 1 to 14 is applied to the soil where plants grow or are capable of growing, to the leaves and/or the fruits of plants or to the seeds of plants.
- 15 16. Method according to Claim 15, characterized in that the fungicidal composition is applied by spraying a liquid to the aerial parts of the crops to be treated.
- 20 17. Method according to either of Claims 15 and 16, characterized in that the quantity of fungicidal composition corresponds to a dose of compound (I) and of compound (II) of between about 0.1 g/ha and about 1 000 g/ha.
- 25 18. Method according to one of Claims 15 to 17, characterized in that the crop treated is chosen from grapevine, cereal crops (wheat, barley, maize, rice) and vegetable crops (haricot bean, onion, cucurbitaceae, cabbage, potato, tomato, sweet pepper, spinach, pea, lettuce, celery, chicory), fruit crops (strawberry plants, raspberry plants), tree crops (apple trees, pear trees, cherry trees, ginseng, lemon trees, coconut palms, pecan trees, cacao trees, walnut trees, rubber trees, olive trees, poplars, banana trees), sunflower, beetroot, tobacco, hop and ornamental crops.
- 30 19. Product comprising a compound of formula (I) and a compound of formula (II) as a combined preparation for simultaneous, separate or sequential use in controlling the phytopathogenic fungi of crops at a site.

Tammes method diagram (ED50)  
*Plasmodium vivax*




Fig.1

Tamemes method diagram (ED70)  
*Plasmopara viticola*





Fig.2

**Tammes method diagram (ED50)**  
*Phytophthora infestans*



**Fig.3**

Tammes method diagram (ED70)  
*Phytophthora infestans*




Fig.4

Tammes method diagram (ED90)  
*Phytophthora infestans*




Fig. 5

Best Available Copy  
INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 02/13490

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 A01N43/40 A01N47/12		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7 A01N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, CHEM ABS Data		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 99 42447 A (AGREVO UK LIMITED) 26 August 1999 (1999-08-26)	
A	& EP 1 056 723 A 6 December 2000 (2000-12-06) cited in the application	
A	EP 0 775 696 A (KUMIAI CHEMICAL INDUSTRY) 28 May 1997 (1997-05-28) cited in the application	
A	EP 0 472 996 A (BAYER AG) 4 March 1992 (1992-03-04)	
A	EP 0 398 072 A (BAYER AG) 22 November 1990 (1990-11-22)	
<div style="display: flex; justify-content: space-between;"><div><input type="checkbox"/> Further documents are listed in the continuation of box C.</div><div><input checked="" type="checkbox"/> Patent family members are listed in annex.</div></div>		
<div style="display: flex; justify-content: space-between;"><div style="width: 45%;"><p>* Special categories of cited documents:</p><p>*A* document defining the general state of the art which is not considered to be of particular relevance</p><p>*E* earlier document but published on or after the international filing date</p><p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p><p>*O* document referring to an oral disclosure, use, exhibition or other means</p><p>*P* document published prior to the international filing date but later than the priority date claimed</p></div><div style="width: 50%;"><p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p><p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p><p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p><p>*&amp;* document member of the same patent family</p></div></div>		
Date of the actual completion of the international search	Date of mailing of the international search report	
22 January 2003	03/02/2003	
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer  Fort, M

## INTERNATIONAL SEARCH REPORT

Information on patent family members

Internat Application No

PCT/EP 02/13490

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9942447	A	26-08-1999	AU 751032 B2 08-08-2002
			AU 2527199 A 06-09-1999
			BR 9908007 A 30-01-2001
			CA 2319005 A1 26-08-1999
			CN 1291187 T 11-04-2001
			CZ 20002993 A3 14-11-2001
			EP 1056723 A1 06-12-2000
			WO 9942447 A1 26-08-1999
			HU 0100817 A2 30-07-2001
			JP 2002503723 T 05-02-2002
			NO 20004159 A 17-10-2000
			NZ 505954 A 20-12-2002
			PL 342376 A1 04-06-2001
			SI 20356 A 30-04-2001
			SK 12392000 A3 12-03-2001
			TR 200002395 T2 21-11-2000
			TR 200101071 T2 21-06-2002
			US 6503933 B1 07-01-2003
			ZA 9901292 A 13-09-1999
EP 0775696	A	28-05-1997	AU 683383 B2 06-11-1997
			AU 2455695 A 04-03-1996
			BR 9508472 A 28-10-1997
			EP 0775696 A1 28-05-1997
			KR 222107 B1 01-10-1999
			PL 318374 A1 09-06-1997
			RU 2129548 C1 27-04-1999
			US 5789428 A 04-08-1998
			CA 2195064 A1 15-02-1996
			CN 1154694 A ,B 16-07-1997
			ES 2148518 T3 16-10-2000
			HU 76686 A2 28-10-1997
			WO 9604252 A1 15-02-1996
			JP 3127386 B2 22-01-2001
			JP 8176115 A 09-07-1996
			PT 775696 T 29-12-2000
EP 0472996	A	04-03-1992	DE 4026966 A1 27-02-1992
			BR 9103637 A 19-05-1992
			DE 59104468 D1 16-03-1995
			EP 0472996 A1 04-03-1992
			EP 0550788 A1 14-07-1993
			ES 2067813 T3 01-04-1995
			GR 3015191 T3 31-05-1995
			HU 59082 A2 28-04-1992
			IE 912997 A1 26-02-1992
			JP 2091863 C 18-09-1996
			JP 4230652 A 19-08-1992
			JP 7068200 B 26-07-1995
			KR 190298 B1 01-06-1999
			MX 9100720 A1 01-04-1992
			PL 291508 A1 19-10-1992
			PL 294789 A1 17-05-1993
			PT 98696 A ,B 31-07-1992
			RU 2015962 C1 15-07-1994
			US 5453531 A 26-09-1995
			ZA 9106691 A 27-05-1992

Best Available Copy  
INTERNATIONAL SEARCH REPORT  
Information on patent family members

International Application No  
PCT/EP 02/13490

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0398072	A 22-11-1990	DE 3915755 A1	29-11-1990
		BR 9002200 A	13-08-1991
		DD 299638 A5	30-04-1992
		DE 59008394 D1	16-03-1995
		EP 0398072 A2	22-11-1990
		ES 2067589 T3	01-04-1995
		GR 3015610 T3	30-06-1995
		HU 54275 A2	28-02-1991
		IE 65097 B1	04-10-1995
		IL 94348 A	12-04-1994
		JP 2867367 B2	08-03-1999
		JP 3005451 A	11-01-1991
		KR 167336 B1	15-01-1999
		PT 93962 A ,B	08-01-1991
		US 5411987 A	02-05-1995
		US 5516786 A	14-05-1996
		US 5210084 A	11-05-1993
		US 5504109 A	02-04-1996
		ZA 9003605 A	27-02-1991